

Package leaflet: Information for the user
Cetrine Allergy 10 mg Film-coated Tablets

Cetirizine dihydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse

What is in this leaflet:

1. What Cetrine Allergy is and what it is used for
2. What you need to know before you take Cetrine Allergy
3. How to take Cetrine Allergy
4. Possible side effects
5. How to store Cetrine Allergy
6. Contents of the pack and other information

1. What Cetrine Allergy is and what it is used for

The active ingredient of Cetrine Allergy is cetirizine dihydrochloride. Cetrine Allergy is an anti-allergy medicine. It helps control your allergic reaction and its symptoms.

Cetrine Allergy relieves symptoms associated with, for example, hay fever, allergy to dust mites (allergic rhinitis) such as:

- sneezing,
- runny or itchy nose,
- itchy palate,
- itchy, red or tearing eyes.

Cetrine Allergy is also used to relieve the symptoms such as itching and hives (associated with chronic urticaria of unknown cause).

2. What you need to know before you take Cetrine Allergy

Do not take Cetrine Allergy:

- if you are allergic to cetirizine or any of the other ingredients of this medicine (listed in section 6), to hydroxyzine or piperazine derivatives (related active substances of other medicines).
- if you have severe kidney problems

Warnings and precautions:

- Talk to your doctor, pharmacist or nurse before taking Cetrine Allergy if you have **kidney problems**. Your treatment may need to be adjusted.
- If you have predisposition factors of urinary retention (e.g. spinal cord lesions or prostatic hyperplasia)
- If you are an epileptic patient or a patient at risk of convulsions, you should ask your doctor for advice.

- If you are scheduled for allergy testing ask your doctor if you should stop taking Cetrine Allergy for several days before testing. This medicine may affect your allergy test results.

Children and adolescents

Not for children under the age of 6 years

Other medicines and Cetrine Allergy

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Cetrine Allergy with food, drink and alcohol

Please be careful if you drink alcohol during use of this medicine, as it may increase the effects of alcohol.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy: there is limited experience of the use of Cetrine Allergy in pregnant women. You should use this medicine with caution during pregnancy.

Breast-feeding: You should avoid using this medicine if you are breastfeeding an infant, as cetirizine hydrochloride is excreted in human milk.

Driving and using machines

At the recommended dose, Cetrine Allergy is not expected to cause you to be drowsy or less alert. However, if you are a sensitive patient, simultaneous use of alcohol and other nervous depressant agents may additionally affect your ability to drive and use machines.

Cetrine Allergy contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Cetrine Allergy

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Adults and adolescents aged 12 years and over:

The recommended dose is one tablet once a day.

Children aged between 6-12 years: The usual dose is half a tablet twice a day (morning and evening).

These tablets are not intended for children under the age of 6 years.

Patients with kidney problems: if you have moderate kidney problems your doctor may reduce your dose to half a tablet once a day. If you have severe kidney problems your doctor may reduce your dose to half a tablet every 2 days.

Swallow the tablets with a glass of water.
You may take this medicine with or without food.

Duration of treatment:

The duration of the treatment may vary depending on the symptoms.

If you take more Cetrine Allergy than you should

The following symptoms may occur after an overdose of Cetrine Allergy:

Confusion, diarrhoea, dizziness, fatigue, headache, weakness, dilatation of the pupils, itching, restlessness, sedation, drowsiness, unresponsiveness, fast heart rhythm, involuntary trembling and inability to pass water.

If you have accidentally taken more of this medicine than you should please contact your doctor or pharmacist immediately.

If you forget to take Cetrine Allergy

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some very rare side effects could be serious:

If any of the following happens, stop taking this medicine and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Allergic reactions such as sudden difficulty in breathing, speaking and swallowing, swelling of the lips, face and neck, extreme dizziness or collapse, itchy, raised skin rash.
- Unusual bruising or bleeding caused by low blood platelets

Other possible side effects

Tell your doctor if any of the following side effects bother you:

Common: may affect up to 1 in 10 people

headache, dizziness, dry mouth, fatigue, feeling sick, diarrhoea, drowsiness, sore throat, runny nose.

Uncommon: may affect up to 1 in 100 people

stomach upset, extreme fatigue, feeling weak, tingling or numbness in the hands and feet, agitation, itchiness, rash.

Rare: may affect up to 1 in 1,000 people

rapid heart beat, swelling, allergic reactions, changes in liver function, weight increase, convulsions, aggression, confusion, depression, hallucination, difficulty in sleeping, red and/or blotchy skin rash.

Very rare: may affect up to 1 in 10,000 people

difficulty focusing, blurred vision, unusual eye movements, fainting, tremor, altered taste, tics, pain and/or difficulty passing water, fixed drug eruption.

Not known: frequency cannot be estimated from the available data

Increase appetite, impairment or loss of memory, vertigo, urinary retention

In very rare cases people have thought about committing suicide, if you feel this way stop taking the tablets and see your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPR

Pharmacovigilance, Earlsfort Terrace, IRL-Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517.

Website: www.hpra.ie; e-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Cetrine Allergy

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP.

The expiry date refers to the last day of that month.

Do not store above 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Cetrine Allergy contains:

- The active substance is cetirizine dihydrochloride. Each film-coated tablet contains 10 mg cetirizine dihydrochloride.
- The other ingredients are: lactose monohydrate, microcrystalline cellulose, colloidal anhydrous silica, magnesium stearate, hypromellose, titanium dioxide (E171) and macrogol 4000.

What Cetrine Allergy looks like and contents of the pack:

Cetrine Allergy are white, oblong film-coated tablets with a score line on one side

PVC/Aluminium blister: 7, 10, 28 and 30 film-coated tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

Rowex Ltd., Bantry, Co. Cork, Ireland

Manufacturer:

Salutas Pharma GmbH, Otto-von-Guericke Allee 1, 39179 Barleben, Germany

Rowa Pharmaceuticals Ltd., Bantry, Co. Cork.

This leaflet was last revised in 07/2014