

The Panadol logo is a stylized white arrow pointing right, containing the word "Panadol" in a bold, white, sans-serif font. Below the arrow is the word "ActiFast" in a white, italicized, sans-serif font.

Panadol

ActiFast

500mg Tablets Paracetamol

PATIENT INFORMATION LEAFLET

Please read right through this leaflet before you start using this medicine.

This medicine is available without prescription, but you still need to use Panadol ActiFast tablets carefully to get the best results from them.

- Keep this leaflet, you may need to read it again.
- If you have any further questions, ask your pharmacist.

In this leaflet:

1. What Panadol ActiFast tablets are and what they are used for
2. Check before you take Panadol ActiFast tablets
3. How to take Panadol ActiFast tablets
4. Possible side effects
5. How to store and dispose of Panadol ActiFast tablets
6. Further information

1. What Panadol ActiFast tablets are and what they are used for

Panadol ActiFast tablets are used for fast pain relief of headaches, toothache, backache, rheumatic and muscle pains and period pain. It also relieves sore throat and the feverishness, aches and pains of colds and flu.

The active ingredient is paracetamol which is a painkiller and also reduces your

temperature when you have a fever. The tablets have been specially formulated so that the paracetamol is absorbed into the bloodstream twice as fast as standard paracetamol tablets and so gets to the source of pain fast.

2. Check before you take Panadol ActiFast tablets



Do not take Panadol ActiFast tablets:

- if you have ever had an **allergic reaction** to paracetamol or to any of the other ingredients (listed in Section 6).
- if you are **taking other medicines containing paracetamol**.
- if you are **under 12 years** unless your doctor tells you to.



Ask your doctor before you take this medicine:

- if you have **severe liver or kidney disease**, including **alcoholic liver disease**.
- if you are on a **sodium-controlled diet**. Each tablet contains **173mg of sodium**.



If you are taking other medicines

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly **metoclopramide** or **domperidone** (for **nausea** [feeling sick] or **vomiting** [being sick]); **colestyramine** (to lower **blood cholesterol**). If you take **blood thinning drugs (anticoagulants e.g. warfarin)** and you need to take a pain reliever on a daily basis, **talk to your doctor** because of the risk of bleeding. However you can still take **occasional** doses of Panadol ActiFast tablets at the same time as anticoagulants.



Pregnancy and breastfeeding

Talk to your doctor before taking Panadol ActiFast tablets if you are **pregnant**. You can take this product whilst **breastfeeding**.

3. How to take Panadol ActiFast tablets



Adults, including the elderly, and children aged 12 years and over:

Swallow 1-2 tablets with half a tumbler of water (100ml), every 4-6 hours as needed.



- **Do not take for more than 3 days without consulting your doctor.**
- **Do not take more frequently than every 4 hours.**
- **Do not take more than 8 tablets in 24 hours.**
- **Do not take more than the recommended dose.**
- **If symptoms persist consult your doctor.**

If you take too many tablets

Immediate medical advice should be sought in the event of an overdose, even if you feel well, because of the risk of delayed, serious liver damage.

4. Possible side effects

Like all medicines Panadol ActiFast tablets can have side effects, but not everybody gets them:

Stop taking this medicine and tell your doctor immediately if:

- you experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- you experience a severe skin rash or peeling of the skin which may be accompanied by mouth ulcers.
- you may have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction to this product.
- you experience unexplained bruising or bleeding.

These reactions are rare.

If you do get any side effects, even those not mentioned in this leaflet, tell your doctor or pharmacist.

5. How to store and dispose of Panadol ActiFast tablets

Keep out of the reach and sight of children.

No special storage conditions are required.

Do not use this medicine after the expiry date (EXP) which is shown on the outer carton and the blisters. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

Active ingredient Each tablet contains Paracetamol 500mg.

Other ingredients Sodium hydrogen carbonate, tricalcium phosphate, starch pregelatinised, povidone, maize starch, potassium sorbate (E 202), microcrystalline cellulose, magnesium stearate, carnauba wax, titanium dioxide (E 171), polydextrose, hypromellose, triacetin and macrogol.

Panadol ActiFast tablets are white, capsule shaped, film-coated tablets and are marked with 'p' on one face.

Panadol ActiFast tablets are produced in packs of 4, 6, 8, 10, 12, 16, 20, 24 or 32 tablets. Not all pack sizes may be marketed.

The Product Authorisation holder is GlaxoSmithKline Consumer Healthcare (Ireland) Limited, Stonemasons Way, Rathfarnham, Dublin 16 and all enquires should be sent to this address.

The manufacturer is GlaxoSmithKline Dungarvan Ltd., Knockbrack, Dungarvan, Co. Waterford, Ireland.

This leaflet was last revised in November 2010.

Panadol and **ActiFast** are registered trade marks of the GlaxoSmithKline group of companies.

PA 678/39/12

